

## ARTICLE

# Alpha 1 Antitrypsin Deficiency Alleles Are Associated With Joint Dislocation and Scoliosis in Williams Syndrome

COLLEEN A. MORRIS,\* ARIEL M. PANI, CAROLYN B. MERVIS, CECILIA M. RIOS, DORIS J. KISTLER, AND RONALD G. GREGG

Elastin haploinsufficiency is responsible for a significant portion of the Williams syndrome (WS) phenotype including hoarse voice, supravalvular aortic stenosis (SVAS), hernias, diverticuli of bowel and bladder, soft skin, and joint abnormalities. All of the connective tissue signs and symptoms are variable in the WS population, but few factors other than age and gender are known to influence the phenotype. We examined a cohort of 205 individuals with WS for mutations in *SERPINA1*, the gene that encodes alpha-1-antitrypsin (AAT), the inhibitor of elastase. Individuals with classic WS deletions and *SERPINA1* genotypes PIMS or PIMZ were more likely than those with a *SERPINA1* PIMM genotype to have joint dislocation or scoliosis. However, carrier status for AAT deficiency was not correlated with presence of inguinal hernia or with presence or severity of SVAS. These findings suggest that genes important in elastin metabolism are candidates for variability in the connective tissue abnormalities in WS. © 2010 Wiley-Liss, Inc.

**KEY WORDS:** Williams syndrome; elastin; alpha-1-antitrypsin; *SERPINA1*; scoliosis; joint dislocation; supravalvular aortic stenosis; fibrillin-1; elastic fibers

**How to cite this article:** Morris CA, Pani AM, Mervis CB, Rios CM, Kistler DJ, Gregg RG. 2010. Alpha 1 antitrypsin deficiency alleles are associated with joint dislocation and scoliosis in Williams syndrome. *Am J Med Genet Part C Semin Med Genet* 154C:299–306.

## INTRODUCTION

Williams syndrome (WS) is a multi-system disorder caused by deletion of 26 contiguous genes on chromosome 7q11.23. Haploinsufficiency for the elastin gene in this region accounts for many of the signs and symptoms of the disorder, including the most significant

cause of morbidity and mortality, supravalvular aortic stenosis (SVAS). Other phenotypic features of WS related to decreased elastin include hoarse voice, hernias, bowel, and bladder diverticuli, and joint abnormalities. All of these phenotypic features have variable severity among individuals with WS. Some of the variability is accounted for

by age; for example, bowel and bladder diverticuli are most commonly observed in adults [Morris et al., 1990]. Some symptoms vary by gender; for example, severe SVAS is more common in males than in females [Sadler et al., 2001]. However, these factors do not explain all of the phenotypic variation. To examine the potential modifying relation of other

Colleen A. Morris, M.D. is Chief of the Genetics Division in the Department of Pediatrics at the University of Nevada School of Medicine. She has studied the natural history and genotype–phenotype correlations of Williams syndrome for 25 years and also has research interests in syndrome delineation and identification and treatment of children with Fetal Alcohol Spectrum Disorders.

Ariel M. Pani, B.A., is a doctoral candidate in the Committee on Evolutionary Biology at the University of Chicago where his primary research focus is on the evolutionary origins of vertebrate CNS signaling centers. He has previously conducted research on genotype/phenotype relations in Williams syndrome.

Carolyn B. Mervis, Ph.D. is a Distinguished University Scholar and Professor of Psychological and Brain Sciences at the University of Louisville. Her primary research focus is on the language, cognitive, social-emotional, and behavioral development of children with Williams syndrome, duplication of the Williams syndrome region, and Down syndrome. She also conducts research on neuroimaging and genotype/phenotype correlations involving the Williams syndrome region.

Cecilia Rios, B.S., M.B.A. is a cytogenetic technologist in the Molecular Cytogenetics Laboratory, Department of Pediatrics at the University of Nevada School of Medicine. Her research focus is the development of FISH probes for characterization of unique deletions, duplications, and inversions.

Doris J. Kistler, Ph.D. is a Research Professor of Psychological and Brain Sciences at the University of Louisville. Her primary research focus is on the development of auditory processing skills and spatial hearing in children with normal hearing and with hearing disorders.

Ronald G. Gregg, Ph.D. is a University Scholar, and Professor and Chair of Biochemistry and Molecular Biology at the University of Louisville. His primary research focus is the use of molecular biology and genetics to study disease. His studies focus on the molecular basis of genome alterations that cause Williams syndrome. He also studies genes that cause congenital stationary night blindness, with the goal of identifying signal transduction pathways in subsets of retinal neurons.

Grant sponsor: National Institute of Neurological Disorders and Stroke; Grant number: R01 NS35102.

\*Correspondence to: Colleen A. Morris, M.D., 2040 W Charleston Blvd, Ste 401, Las Vegas, NV 89102. E-mail: cmorris@medicine.nevada.edu  
DOI 10.1002/ajmg.c.30265

Published online 21 April 2010 in Wiley InterScience (www.interscience.wiley.com)

© 2010 Wiley-Liss, Inc.

# Alpha Associated 1 Antitrypsin With Joint Deficiency Dislocation Alleles and Scoliosis Are in Williams Syndrome

COLLEEN A. MORRIS,\* ARIEL M. PANI, CAROLYN B. MERVIS, CECILIA M. RIOS, DORIS J. KISTLER,

AND

RONALD G. GREGG

Elastin haploinsufficiency is responsible for a significant portion of the Williams syndrome (WS) phenotype including hoarse voice, supra-valvar aortic stenosis (SVAS), hernias, diverticuli of bowel and bladder, soft skin, and joint abnormalities. All of the connective tissue signs and symptoms are variable in the WS population, but few factors other than age and gender are known to influence the phenotype. We examined a cohort of 205 individuals with WS for mutations in *SERPINA1*, the gene that encodes alpha-1-antitrypsin (AAT), the inhibitor of elastase. Individuals with classic WS deletions and *SERPINA1* genotypes PiMS or PiMZ were more likely than those with a *SERPINA1* PiMM genotype to have joint dislocation or scoliosis. However, carrier status for AAT deficiency was not correlated with presence of inguinal hernia or with presence or severity of SVAS. These findings suggest that genes important in elastin metabolism are candidates for variability in the connective tissue abnormalities in WS. © 2010 Wiley-Liss, Inc.

**KEY WORDS:** Williams syndrome; elastin; alpha-1-antitrypsin; *SERPINA1*; scoliosis; joint dislocation; supra-valvar aortic stenosis; fibrillin-1; elastic fibers

How to cite this article: Morris CA, Pani AM, Mervis CB, Rios CM, Kistler DJ, Gregg RG. 2010. Alpha 1 antitrypsin deficiency alleles are associated with joint dislocation and scoliosis in Williams syndrome. *Am J Med Genet Part C Semin Med Genet* 154C:299–306.

## INTRODUCTION

Williams syndrome (WS) is a multi-system disorder caused by deletion of 26 contiguous genes on chromosome 7q11.23. Haploinsufficiency for the elastin gene in this region accounts for many of the signs and symptoms of the disorder, including the most significant

cause of morbidity and mortality, supra-valvar aortic stenosis (SVAS). Other phenotypic features of WS related to decreased elastin include hoarse voice, hernias, bowel, and bladder diverticuli, and joint abnormalities. All of these phenotypic features have variable severity among individuals with WS. Some of the variability is accounted for

by age; for example, bowel and bladder diverticuli are most commonly observed in adults [Morris et al., 1990]. Some symptoms vary by gender; for example, severe SVAS is more common in males than in females [Sadler et al., 2001]. However, these factors do not explain all of the phenotypic variation. To examine the potential modifying relation of other

Colleen A. Morris, M.D. is Chief of the Genetics Division in the Department of Pediatrics at the University of Nevada School of Medicine. She has studied the natural history and genotype–phenotype correlations of Williams syndrome for 25 years and also has research interests in syndrome delineation and identification and treatment of children with Fetal Alcohol Spectrum Disorders.

Ariel M. Pani, B.A., is a doctoral candidate in the Committee on Evolutionary Biology at the University of Chicago where his primary research focus is on the evolutionary origins of vertebrate CNS signaling centers. He has previously conducted research on genotype/phenotype relations in Williams syndrome.

Carolyn B. Mervis, Ph.D. is a Distinguished University Scholar and Professor of Psychological and Brain Sciences at the University of Louisville. Her primary research focus is on the language, cognitive, social-emotional, and behavioral development of children with Williams syndrome, duplication of the Williams syndrome region, and Down syndrome. She also conducts research on neuroimaging and genotype/phenotype correlations involving the Williams syndrome region.

Cecilia Rios, B.S., M.B.A. is a cytogenetic technologist in the Molecular Cytogenetics Laboratory, Department of Pediatrics at the University of Nevada School of Medicine. Her research focus is the development of FISH probes for characterization of unique deletions, duplications, and inversions.

Doris J. Kistler, Ph.D. is a Research Professor of Psychological and Brain Sciences at the University of Louisville. Her primary research focus is on the development of auditory processing skills and spatial hearing in children with normal hearing and with hearing disorders.

Ronald G. Gregg, Ph.D. is a University Scholar, and Professor and Chair of Biochemistry and Molecular Biology at the University of Louisville. His primary research focus is the use of molecular biology and genetics to study disease. His studies focus on the molecular basis of genome alterations that cause Williams syndrome. He also studies genes that cause congenital stationary night blindness, with the goal of identifying signal transduction pathways in subsets of retinal neurons.

Grant sponsor: National Institute of Neurological Disorders and Stroke; Grant number: R01 NS35102.

\*Correspondence to: Colleen A. Morris, M.D., 2040 W Charleston Blvd. Ste 401, Las Vegas, NV 89102. E-mail: cmorris@medicine.nevada.edu DOI 10.1002/ajmg.c.30265 Published online 21 April 2010 in Wiley InterScience (www.interscience.wiley.com)

© 2010 Wiley-Liss, Inc.

genes to the connective tissue and cardiovascular phenotypes in WS, we elected to examine a common genotype known to be associated with variation in elastin metabolism. Alpha-1 antitrypsin (AAT) inhibits elastase; deficiency of this protein is associated with emphysema and liver disease in homozygotes, but heterozygotes also have lower than normal levels of AAT that can adversely impact health. Mutant alleles of the causative gene, SERPINA1, are common in the general population, so we reasoned that our cohort of participants with WS would include heterozygotes. The purpose of the study was to determine if mutations in SERPINA1 influenced the connective tissue phenotypes in WS.

Elastin deficiency is the proximal cause for the connective tissue abnormalities in WS, but other factors likely influence phenotypic variability. Elastin is a major component of elastic fibers that confer elasticity and recoil to several tissues. The organization of elastic fibers depends on the tissue. Elastic fiber networks are found in hollow organs such as the lung, bowel, and bladder, whereas linear arrangements of elastic fibers in association with collagen are typical of ligaments and skin. Concentric rings of elastin are present in arteries and intervertebral discs. In arteries, the elastic fiber rings are interspersed with vascular smooth muscle cells; each ring of elastic fibers and muscle cells forms a lamellar unit. Most elastin is produced in the third trimester of pregnancy and in the first year of life. Turnover is low, but biomechanical stress and resultant inflammation can expose elastic fibers to degradation. Elastase is the primary enzyme that causes elastin degradation.

Alpha 1 antitrypsin (AAT) is a glycoprotein that inhibits elastase, and its deficiency increases the risk for development of chronic obstructive pulmonary disease (COPD) and chronic liver disease [Zorzetto et al., 2008]. The genotype for the SERPINA1 gene determines AAT levels in the serum. The normal gene is labeled PiM and the common deficiency alleles are PiS, which expresses 50% of normal AAT, and PiZ, which expresses 20% [De

300 AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) ARTICLE

Serres et al., 2003]. The lowest levels of AAT are found with PiZZ and PiSZ genotypes, but diminished AAT levels have been demonstrated in populations with PiMS and PiMZ genotypes [Zorzetto et al., 2008]. The genotype frequencies vary by population; in the USA, the PiMS frequency is 5.88% and PiMZ is 2.78% [De Serres et al., 2003]. Low levels of AAT result in decreased elastase inhibition and therefore increased elastin degradation with inflammation.

### **Low levels of AAT result in decreased elastase inhibition and therefore increased elastin degradation with inflammation.**

There are many connective tissue abnormalities in WS, affecting multiple organ systems. Some signs are nearly universal in WS, such as the hoarse voice that is present in 98% and is due to decreased elastic fibers in the lamina propria of the vocal folds [Hammond et al., 1998; Vaux et al., 2003; Watts et al., 2008]. Other traits present in over 90% include periorbital fullness and full cheeks in infants, lax joints in infancy, and soft skin [Morris et al., 1988]. While joints are loose in infancy, joint contractures typically begin to appear in childhood, worsen with age, and most predominantly affect the lower extremities [Morris et al., 1988, 1990; Kaplan et al., 1989]. The combination of central hypotonia and lax joints in young children contributes to delayed walking. The young child will often develop abnormal compensatory postures, including mild lordosis, and kyphosis. The above-mentioned clinical features are present in most individuals with WS, though some vary with age. Elastin deficiency alone is sufficient explanation for occurrence of these traits. Unlike lordosis and kyphosis, however, scoliosis is a fairly low frequency finding in WS, and therefore was one of the traits targeted for investigation in the

current study. In a previous study of a cohort of 42 individuals aged 1–34 years with WS, 12% had scoliosis, 38% had inguinal hernia, and 64% had SVAS [Morris et al., 1988].

Inguinal hernias are typically diagnosed in infancy in WS and occur in 40% [Morris, in press] as compared with 5% incidence in the general pediatric population [Brandt, 2008]. Because decreased elastic fibers in the abdominal wall as a consequence of aging are associated with inguinal hernias in the general population, we investigated the possibility that AAT deficiency might be associated with hernia occurrence in WS. Bowel and bladder diverticuli

are common in WS, but usually present in adulthood, although at an earlier age than in the general population [Poher and Morris, 2007]. Because diverticuli usually are not detected until symptoms warrant invasive investigation, the prevalence in the WS population is unknown and therefore could not be evaluated in this study. The most serious consequence of the ELN deletion in WS is elastin arteriopathy. Aortic smooth muscle cells from individuals with WS produce low levels of elastin mRNA and thus deposit less elastin protein in tissues, resulting in sparse elastic fibers, which in turn results in increased proliferation of smooth muscle cells [Urban et al., 2002].

The most serious consequence of the ELN deletion in WS is elastin arteriopathy. Aortic smooth muscle cells from individuals with WS produce low levels of elastin mRNA and thus deposit less elastin protein in tissues, resulting in sparse elastic fibers, which in turn results in increased proliferation of smooth muscle cells.

In the ascending aorta of WS, the lamellar units are increased from the

normal number of 40 to 120, resulting in thickening of the media of the vessel [Dridi et al., 2005]. This change in the number of lamellar units is present at birth [Wagenseil and Mecham, 2009], indicating the ELN mutation is the etiology of the abnormality of the media. However, there can be sites of discrete narrowing that occur in areas of hemodynamic stress and turbulence. The proliferation of arterial smooth muscle cells that occurs secondary to elastin deficiency leads to a progression of pathology in that hyperplastic intimal lesions develop [Urban et al., 2002], encroaching on the lumen of the artery. Since these areas would be subject to tissue injury, we hypothesized that the severity of the SVAS may be related to elastin degradation. If so, then lowered AAT levels could be associated with a more severe arterial phenotype. Previous studies have shown that approximately 30% of individuals with SVAS require surgical intervention [Kececioglu et al., 1993].

## METHODS

### Participants and Clinical Assessment

The study population included 205 individuals with WS (107 females, 98 males), all of whom had the typical deletion on chromosome 7q11.23. Informed consent and specimens were obtained following IRB-approved protocols. Medical records were reviewed on all participants, and physical examinations were performed on 174/205 participants by C.A.M. Age at the time of examination ranged from 9 months to 45 years. Inguinal hernia and joint dislocation were scored as present or absent. Scoliosis was assessed by physical inspection including the Adams forward bending test and review of radiography reports. Curves greater than 10 degrees were considered positive for scoliosis. Individuals were coded as negative for scoliosis if they had no physical/radiographic evidence of scoliosis by age 8 years. Children <8 years old were excluded from analysis because they could still develop spinal curvature.

### ARTICLE AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) 301

None of the participants had vertebral malformations. For individuals who had not had surgery for SVAS, cardiovascular disease severity was scored based on review of echocardiography reports using the recorded Doppler measurements (Table I). Individuals who required surgical correction for SVAS were classified as severely affected.

### Genotype Analysis

Cytogenetic preparations were made from either fresh whole blood or previously established lymphoblastoid lines, and standard methods for metaphase FISH experiments to establish the size of deletion were followed as previously described [Hobart et al., in press, this volume]. All participants had typical deletions of 1.55–1.8Mb of chromosome 7q11.23.

Genomic DNA was isolated from lymphoblastoid cell lines or whole blood using a Puregene DNA isolation kit (Gentra Systems, Minneapolis, MN) and genotyping was performed by pyrosequencing analysis. The SERPINA1 Pi Z and Pi S loci were amplified by PCR using the forward primers PiZ 50-TTGACCTCGGGGGGATAG-30 and PiS 50-TTGGTGATGATATCGTG-GGTGAGT-30 and reverse primers PiZ 50(bio)-GGGATCAGCCTTACA-ACGTGTC-30 and PiS 50(bio)-CAAT-GCCACCGCCATCTT-30. For each individual, 5 µl of biotinylated PCR product was analyzed on a Biotage PSQ HS 96 pyrosequencer (Biotage, Uppsala, Sweden) using sequencing primers PiZ 50-AGCTTCAGTCC-CTTTCT-30 and PiS 50-CGTGGGT-GAGTTCATTT-30. Genotypes were called by the instrument software in

TABLE I. Classification for SVAS Severity Based on Echocardiography Results

Degree of severity of SVAS Maximum velocity (m/sec) Peak gradient (mm/Hg)

None <2 <10 Mild 2–2.5 11–30 Moderate 2.6–3.5 31–60 Severe >3.5 >60

SNP analysis mode, and pyrograms were visually checked for errors.

## RESULTS

## Clinical Assessment

We identified four individuals with WS who had joint dislocations, a finding not previously reported in WS. All four individuals in this cohort were heterozygous for AAT mutations. An 18-month-old girl, genotype PiMZ, had dislocation of the left shoulder. A 3.7-year-old girl, genotype PiMS, had bilateral dislocated knees treated surgically. A boy, genotype PiMZ, had a dislocated knee at age 7 years and also was noted to have mild scoliosis. A male, genotype PiMZ, with a history of mild scoliosis had a dislocated right knee treated surgically at age 16 years.

The prevalence of scoliosis in the participants aged 8 years and older was 18%. Most children had a juvenile (between ages 5 and 10 years) presentation of the scoliosis. For some adults in the group, all with mild untreated scoliosis at evaluation, age of onset of scoliosis was not documented. There was no significant difference in the rates of scoliosis between males (13/55) and females (7/56) (Fisher's exact test,  $P = 1/40.145$ ). Severe scoliosis requiring bracing or surgery was present in six participants (Fig. 1).

Inguinal hernia (indirect) was diagnosed in infancy in 26% of the participants. A significantly higher proportion of males (38/98) than females (17/107) was affected (Fisher's exact test,  $P < 0.0001$ ).

SVAS was documented in 68%; of those affected, 31.4% required surgical correction. The difference in proportion

of males (72/98) and females (68/107) who had any SVAS was not significant (Fisher's exact test,  $P=0.136$ ). However, a significantly larger proportion of males (29/97) than females (15/107) had severe SVAS (Fisher's exact test,  $P=0.007$ ).

#### 302 AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) ARTICLE

Figure 1. Spine radiograph of a 12-year-old female with WS and complex scoliosis. The onset of scoliosis was age 6 years, and she currently is treated with a brace. Her AAT genotype is PiMS.

#### Genotyping

All participants had the typical WS deletion of 7q11.23 documented by FISH. Observed gene frequencies for each of the possible SERPINA1

genotypes are reported in Table II. The observed percentage for the MZ genotype was 4.88%. We calculated the 95% confidence interval for this percentage using the modified Wald method [Agresti and Couli, 1998]. The 95% confidence interval (2.56–8.85%) included the estimate of 2.78% for the USA previously reported by De Serres et al. [2003] for this genotype. For the MS genotype, the observed percentage was 4.40%. The estimate of 5.88% reported by De Serres et al. [2003] for this genotype is included in the 95% confidence interval for the observed percentage (2.21–8.25%). Thus, the observed percentages of the MZ and MS genotypes for the present cohort are consistent with previous estimates for the USA. Genotypes were in Hardy–Weinberg equilibrium, and there were no PiSZ compound heterozygotes observed.

#### Genotype–Phenotype Correlation

We examined the association between SERPINA1 genotype and a diagnosis of joint dislocation, inguinal hernia, any SVAS, or severe SVAS for 205 individuals with WS using Fisher's exact tests (Table III). Due to the low prevalence of MZ and MS genotypes, they were combined into a single category of AAT deficiency carriers for analyses. Because the onset of scoliosis is variable but typically appears in WS prior to age 8 years, only individuals 8 years or older were included in the analysis for scoliosis ( $n=111$ ). We found that AAT deficiency carriers with WS were significantly more likely than non-carriers to have a diagnosis of joint dislocation ( $P<0.0001$ ) or scoliosis ( $P<0.0001$ ).

AAT deficiency carriers with WS were not more likely than non-carriers to have inguinal hernia ( $P=0.171$ ). However, in contrast to the sex difference found for the full sample, there was no difference in the rate of inguinal hernia for male (4/10) or female (4/9) AAT deficiency carriers (Fisher's exact test,  $P=1.00$ ).

AAT deficiency carriers also were not more likely than non-carriers to have either any SVAS ( $P=1.00$ ), or



severe SVAS (P1/40.231). However, in contrast to the significant sex difference in proportion of individuals with severe SVAS that was found for the full sample, the difference in proportion of male (4/9) and female (2/9) AAT deficiency carriers who had severe SVAS was not significant (Fisher's exact test, P1/40.620).

## DISCUSSION

As is true with all syndromes, each aspect of the WS phenotype is variable. The connective tissue phenotype is known to be due to decreased elastin production, resulting in diminished elastic fibers in all tissues studied. For traits that are nearly universal in WS, paucity of elastic fibers is sufficient to explain the phenotype. For those features that have a more varied presentation or wide range of severity, other factors are likely contributing to the phenotype.

### Elastin Arteriopathy

No effect of SERPINA1 mutation status was found for the severity of SVAS. While the media of the arterial wall of most medium and large arteries in WS is thick compared to normal, the resultant

### ARTICLE AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) 303

TABLE II. Allele and Genotype Frequencies of SERPINA1 PiZ and PiS in Williams Syndrome

Allele frequency Genotype frequency

Pi Z PiS

M Z	M S	MM	MZ	ZZ	MM	MS	SS	Proportion
400/410	10/410	401/410	9/410	195/205	10/205	0/205	196/205	9/205
97.56%	2.44%	97.80%	2.20%	95.12%	4.88%	0.00%	95.60%	4.40%
0.00%								0.00%

TABLE III. Association of a-1-Antitrypsin Genotype With Connective Tissue Pathologies in Williams Syndrome

Diagnosis

luminal narrowing reaches clinical sig-

decreased

occur

renal

aortic

outflow

arteries

most

SVAS

SVAS

severe SVAS and male gender, in agreement only if there is

ment with the cross-sectional study of perfusion of an organ, such as may

Sadler et al. [2001]. However, gender in a localized narrowing of the

was not found to be associated with artery; or if narrowing above the

SVAS severity in the recent study of or pulmonary valve results in

individuals with WS evaluated by a tract obstruction that leads to cardiac cardiologist before age one year [Collins hypertrophy. Narrowing of the

et al., 2010]. The variability of the SVAS in the pulmonary circulation

phenotype cannot be explained by typically improves over time, but

gender alone. may worsen, most commonly in the first

While elastase is the primary 5 years of life [Collins et al., 2010].

about 30%

requiring

demonstrated in

previously

al., 1993;

al.,

between

MMPs in aortic tissue samples from WS, isolated SVAS, and controls and found SVAS severity is quite variable, with about 30% of individuals

increased staining for MMP2, MMP7, and MMP9 in WS and isolated SVAS compared to controls. However, who have SVAS requiring surgical treatment, as

the tissue samples from affected individuals were removed during aortic surgery, so the individuals had a severe cardiovascular phenotype demonstrated in this study as well as in previously reported series.

vascular phenotype. Therefore, the increased presence of the MMPs could be primary or secondary to the aortic wall disease. Further investigation of connective tissue physiology in WS and

$\alpha$ -1-Antitrypsin heterozygotes (MZ or MS) Normal (MM)

Prevalence Percentage affected n Pa Joint dislocation 4/19 21% 0/186  
00% 205 <0.0001 Scoliosis 8/11 73% 12/100 12% 111 <0.0001 Inguinal hernia 8/19 42% 47/186 25% 205 0.171 Any SVAS 13/  
19 68% 127/186 67% 205 1.000 Severe SVAS 6/18 33% 38/186 20% 204 0.231

aFrom Fisher's exact test.

enzyme responsible for elastin degradation, severity is quite variable, with  
tion, metalloproteinases (MMPs) are also of individuals who have SVAS  
able to degrade elastin. MMPs increase surgical treatment, as  
in tissues in response to inflammation or this study as well as in  
injury. To investigate the possibility that reported series [Kececioglu et  
MMPs are involved in the WS arterio- Del Pasqua et al., 2009; Collins et  
pathy, Dridi et al. [2005] immunolabeled 2010]. We found an association

in animal models may provide additional clues into the basis of phenotypic variability.

### Inguinal Hernia

Inguinal hernia is more common in WS (38% in a previous series [Morris et al., 1988] and 26% in the current study) than in the general population (5% [Brandt, 2008]). We found that males with WS were more commonly affected; in the general population the male to female ratio is 10:1 [Brandt, 2008]. Heterozygous mutation in SERPINA1 was not associated with indirect inguinal hernia in individuals with WS in the current study. Inguinal hernia in adults from the general population is associated with decreased elastin synthesis and increased enzymatic degradation of elastin in the transversalis fascia of individuals with direct inguinal hernia, but not indirect inguinal hernia [Pascual et al., 2009]. This finding suggests that AAT mutations could play a role in direct inguinal hernia; however, none of the individuals in our study had this form of hernia.

### Scoliosis

Scoliosis, a lateral curvature of the spine, is classified as congenital (most often associated with malformations of the spine such as hemivertebrae), neuromuscular (associated with myopathies, cerebral palsy, etc.), syndrome-related (e.g., Marfan syndrome (MFS), neurofibromatosis type 1), or idiopathic, which is the most common type. Idiopathic scoliosis is further subdivided by age of onset: infantile, 0–4 years, accounting for 0.5%; juvenile, 4–10 years, 10.5%; and adolescent, >10 years, 89% [Janicki and Alman, 2007]. Most individuals with idiopathic scoliosis have mild curvature that does not require treatment. Although 2.5% of the general population has adolescent idiopathic scoliosis (AIS), only 0.25% of the population requires surgical intervention. The female to male ratio in the surgically treated group is 7–1 [Asher and Burton, 2006]. Progression of the curve is most likely to occur during periods of rapid growth.

### 304 AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) ARTICLE

It has long been observed that scoliosis occurs in families [Miller et al., 1996], and it has been hypothesized that genes involved in connective tissue metabolism may contribute to disease if mutations compromise the biomechanical integrity of the connective tissue. Scoliosis occurs commonly with some syndromes of connective tissue, including MFS, which has an incidence of scoliosis of approximately 60% and is caused by mutations in fibrillin 1 (FBN1). Most individuals with scoliosis who have MFS have mild curves, but there are some who have rapid progression. The cause of the variability of the scoliosis phenotype is unknown. In addition, a three generation family with kyphoscoliosis who did not meet the criteria for MFS was found to have a FBN1 mutation [Ades et al., 2002]. FBN1 encodes fibrillin 1, a protein that is one of the two major proteins comprising the elastic fiber. The other protein is elastin, which is deficient in WS. Because scoliosis is more common in MFS than in WS, it appears that FBN1 mutation confers a greater risk for spinal curvature, perhaps due to a dominant negative effect. An alternative explanation is that the unique mechanical properties of elastin and fibrillin 1 account for the differing prevalence of scoliosis in the two syndromes. In studies of the vascular mechanics in mice haploinsufficient for both ELN and FBN1, Carta et al. [2009] have demonstrated that elastin contributes elastic recoil and fibrillin 1 contributes tensile strength to the aortic wall. If the same principles apply to the intervertebral disc, then fibrillin 1 may have a greater role in accommodating the biomechanical stress occurring during periods of rapid growth.

The intervertebral disc is well organized: Yu et al. [2005] demonstrated that the human annulus fibrosus consists of collagen fiber bundles organized in concentric lamellae alternating with dense elastic fibers around the nucleus pulposus. Elastic fibers are also present within the lamellae, parallel with the collagen bundles. The lamellae of the inner annulus have a denser concentration of elastic fibers that are further

organized, oriented at 60 or 120 degree angles to those fibers in the adjacent lamella. This well ordered architecture is severely disrupted in scoliotic intervertebral discs, and elastic fibers are sparse [Yu et al., 2005].

During periods of rapid growth, the spine and its joints—the intervertebral discs—are subject to mechanical force. Tissue injury is associated with increased levels of elastase, which degrades elastin in the extracellular matrix. The findings of the current study indicate that individuals with WS who were heterozygotes for AAT deficiency,

and accordingly would have lower levels of AAT and therefore impaired inhibition of elastase, were more likely to develop scoliosis. A similar effect for a

**The findings of the current study indicate that individuals with WS who were heterozygotes for AAT deficiency, and accordingly would have lower levels of AAT and therefore impaired inhibition of elastase, were more likely to develop scoliosis.**

different genetic polymorphism has been demonstrated in a case control study of scoliosis in the general population. Aulisa et al. [2007] found a higher risk of scoliosis in individuals who had either polymorphisms in the matrix metalloproteinase-3 (MMP-3) gene or polymorphism of the promoter region of the interleukin-6 (IL-6) gene. Both genes are involved in inflammation. The 5A/5A genotype of MMP-3 conferred a 3.34 odds ratio for scoliosis and the G/G polymorphism of IL-6 resulted in an odds ratio of 10.54 [Aulisa et al., 2007]. Because these polymorphisms are fairly common, studying their associations with severity of scoliosis in populations with abnormalities in extracellular matrix structural proteins, such as MFS and WS, may be fruitful in the

characterization of phenotypic variability in those conditions.

### Joint Dislocation

An unexpected finding was the association of SERPINA1 mutations with joint dislocation, a phenotype that had not previously been reported for individuals with WS. Elastic fibers are a major component of ligaments. In large joints, such as the knee, bundles of elastic fibers also comprise a three dimensional network in the articular capsule. Marked reduction in the numbers of elastic fibers in that tissue has been documented in MFS [Gigante et al., 1999]. Joint dislocation has been reported in MFS; investigation of mutations in AAT in MFS may provide insight into the pathogenesis of joint dislocation in that condition.

### Summary

In summary, the results from this study suggest that AAT mutation carrier status is not a factor in the severity of the elastin arteriopathy in WS. However, it does appear to have an effect on joints/ ligaments. AAT inhibits elastase, which is important in the process of remodeling and repair of the connective tissue. Elastase production increases with inflammation, and an individual with an MZ or MS AAT genotype will have increased degradation of the connective tissue matrix, leading to decreased joint stability. These data add mutations in AAT to the polymorphisms that affect the connective tissue phenotype (scoliosis and joint dislocation) in WS. Future studies should evaluate MMP-3 and IL-6 polymorphisms in WS, as these have been associated with scoliosis in the general population. Finally, the results of this study suggest that mutations in AAT should be evaluated in genotype–phenotype studies of other syndromes associated with scoliosis, such as MFS.

### ACKNOWLEDGMENTS

This study was supported by grant R01 NS35102 from the National Institute of

ARTICLE AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) 305

Neurological Disorders and Stroke. We are grateful for the participation of individuals with Williams syndrome and their families. We thank the series of research coordinators, Stephanie Nelson, Michele Perez, Marisol Gregorio, and Ruth Denton, for arranging medical research trips to collect data on Williams syndrome for the University of Nevada School of Medicine.

### REFERENCES

- Ades LC, Sreetharan D, Onikul E, Stockton V, Watson KC, Holman KJ. 2002. Segregation of a novel FBN1 gene mutation, G1796E, with kyphoscoliosis and radiographic evidence of vertebral dysplasia in three generations. *Am J Med Genet* 109:261–270.
- Agresti A, Couli BA. 1998. Approximate is better than “Exact” for interval estimation of binomial proportions. *Am Stat* 52:119–126.
- Asher MA, Burton DC. 2006. Adolescent idiopathic scoliosis: Natural history and long term treatment effects. *Scoliosis* 1:2.
- Aulisa L, Papaleo P, Pola E, Angelini F, Aulisa AG, Tamburrelli FC, Pola P, Logroscino CA. 2007. Association between IL-6 and MMP-3 gene polymorphisms and adolescent idiopathic scoliosis. *Spine* 32:2700–2702.
- Brandt ML. 2008. Pediatric hernias. *Surg Clin North Am* 88:27–43.
- Carta L, Wagenseil JE, Knutsen RH, Mariko B, Faury G, Davis EC, Starcher B, Mecham RP, Ramirez F. 2009. Discrete contributions of elastic fiber components to arterial development and mechanical compliance. *Arterioscler Thromb Vasc Biol* 29:2083–2089.
- Collins RTII, Kaplan P, Somes GW, Rome JJ. 2010. Cardiovascular abnormalities, interventions, and long term outcomes in infantile Williams syndrome. *J Pediatr* 156: 253–258.
- De Serres FJ, Blanco I, Fernandez-Bustillo E. 2003. Genetic epidemiology of alpha-1-antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand, and the United States of America. *Clin Genet* 64:382–397.
- Del Pasqua A, Rinelli G, Toscano A, Iacobelli R, Digilio C, Marino B, Saffirio C, Mondillo S, Pasquini L, Sanders SP, de Zorzi A. 2009. New findings concerning cardiovascular manifestations emerging from long-term follow-up of 150 patients with the Williams–Beuren syndrome. *Cardiol Young* 19:563–567.
- Dridi SM, Bertaud AF, Tchen SI, Senni K, Ejeil AL, Pellat B, Lyonnet S, Bonnet D, Charpiot P, Godeau G. 2005. Vascular wall remodeling in patients with supravalvular aortic stenosis and Williams Beuren syndrome. *J Vasc Res* 42:190–201.
- Gigante A, Chillemi C, Greco F. 1999. Changes of elastic fibers in musculoskeletal tissues of Marfan syndrome: A possible mechanism of joint laxity and skeletal overgrowth. *J Pediatr Orthop* 19:283–288.
- Hammond TH, Gray SD, Butler J, Zhou R, Hammond E.

1998. Age- and gender- related elastin distribution changes in human vocal folds. *Otolaryngol Head Neck Surg* 119:314–322.

Hobart HH, Morris CA, Mervis CB, Pani AM, Kistler DJ, Rios CM, Kimberley KW, Gregg RG, Bray-Ward P. 2010. Inversion of the Williams syndrome region is a common polymorphism found more frequently in parents of children with Williams syndrome. *Am J Med Genet Part C Semin Med Genet* (in press).

Janicki JA, Alman B. 2007. Scoliosis: Review of diagnosis and treatment. *Paediatr Child Health* 12:771–776.

Kaplan P, Kirschner M, Watters G, Costa T. 1989. Contractures in patients with Williams syndrome. *Pediatrics* 84:771–773.

Kececioglu D, Kotthoff S, Vogt J. 1993. Williams–Beuren syndrome: A 30-year follow-up of natural and postoperative course. *Eur Heart J* 14:1458–1464.

Miller NH, Mims B, Child A, Milewicz DM, Sponseller P, Blanton SH. 1996. Genetic analysis of structural elastic fiber and collagen genes in familial adolescent idiopathic scoliosis. *J Orthop Res* 14:994–999.

Morris CA. 2010. Williams syndrome. In: Cassidy SB, Allanson JE, editors. *Management of genetic syndromes*. 3rd edition. Hoboken, NJ: Wiley.

Morris CA, Dilts C, Demsey SA, Leonard CO, Blackburn B. 1988. The natural history of Williams syndrome: Physical characteristics. *J Pediatr* 113:318–326.

Morris CA, Leonard CO, Dilts C, Demsey SA. 1990. Adults with Williams syndrome. *Am J Med Genet* 6:102–107.

Pascual G, Rodriguez M, Mecham RP, Sommer P, Bujan J, Belion JM. 2009. Lysyl oxidase like-1 dysregulation and its contribution to direct inguinal hernia. *Eur J Clin Invest* 39:328–337.

Pober BR, Morris CA. 2007. Diagnosis and management of medical problems in adults with Williams–Beuren syndrome. *Am J Med Genet Part C* 145C:280–290.

Sadler LS, Pober BR, Grandinetti A, Scheiber D, Fekete G, Sharma AN, Urban Z. 2001. Differences by sex in cardiovascular disease in Williams syndrome. *J Pediatr* 139:849–852.

Urban Z, Riazi S, Seidl TL, Katahira J, Smoot LB, Chitayat D, Boyd CD, Hinek A. 2002. Connection between elastin haploinsufficiency and increased cell proliferation in patients with supravalvular aortic stenosis and Williams–Beuren syndrome. *Am J Hum Genet* 71:30–44.

Vaux KK, Wojtczak H, Benirsche K, Jones KL. 2003. Vocal cord anomalies in Williams syndrome: A further manifestation of elastin deficiency. *Am J Med Genet Part A* 119A:302–304.

Wagenseil JE, Mecham RP. 2009. Vascular extracellular matrix and arterial mechanics. *Physiol Rev* 89:957–989.

Watts CR, Awan SN, Marler JA. 2008. An investigation of voice quality in

individuals with inherited elastin gene abnormalities. *Clin Linguist Phon* 22:199–213. Yu J, Fairbank JCT, Roberts S, Urban JPG. 2005. The elastic fiber network of the annulus

306 AMERICAN JOURNAL OF MEDICAL GENETICS PART C (SEMINARS IN MEDICAL GENETICS) ARTICLE

fibrosus of the normal and scoliotic human intervertebral disc. *Spine* 30:1815–1820. Zorzetto M, Russi E, Senn O, Imboden M, Ferrarotti I, Tinelli C, Campo I, Ottaviani S, von Eckardstein A, Berger W, Brandli O,

Rochat T, Luisetti M, Probst-Hensch N, SAPALDIA Team. 2008. SERPINA1 gene variants in individuals from the general population concentrations. *with Clin reduced Chem* 54:1331–1338.

$\alpha$

1

-antitrypsin